

Contents lists available at [SciVerse ScienceDirect](#)

Biochimica et Biophysica Acta

journal homepage: www.elsevier.com/locate/bbabio

S14 Terminal Oxidases

14L1

Proton transfer in NO-reducing heme-copper oxidases

Pia Ädelroth, Josy ter Beek, Hyun Ju Lee, Nathalie Gonska, Lina Salomonsson, Davinia Arjona, Joachim Reimann
 Stockholm University, Department of Biochemistry and Biophysics,
 Svante Arrhenius väg 16, 10691 Stockholm, Sweden
 E-mail: piaa@dbb.su.se

Heme-copper oxidases (HCuOs) terminate the respiratory chain in mitochondria and most bacteria. They are integral membrane proteins that catalyse the reduction of oxygen, and they convert the liberated free energy into a proton-motive force across the membrane. The HCuO superfamily has been divided into the oxygen-reducing A-, B- and C-type oxidases as well as the bacterial NO reductases (NOR), where the latter are not O₂-reducers but instead catalyse the reduction of NO in the denitrification process. Proton transfer to the catalytic site in the mitochondrial-like A family occurs through two well-defined pathways termed the D- and K-pathways. The B, C, and NOR families differ in the pathways as well as the mechanisms for proton transfer to the active site and across the membrane. In the NORs, recent structural work has shown that there are possible differences in the proton transfer (PT) pathways between the NOR that uses cyt. *c* as electron donor (cNOR) and NOR that uses quinol (qNOR), where qNOR has a putative PT pathway from the cytoplasm, in contrast to cNOR which is known to be non-electrogenic and use periplasmic protons for NO-reduction ($2\text{NO} + 2\text{e}^- + 2\text{H}^+ \rightarrow \text{N}_2\text{O} + \text{H}_2\text{O}$). We study the PT pathways and mechanisms specifically in the C-type HCuO as well as both cNOR and qNOR using site-directed mutagenesis in combination with time-resolved spectroscopy during single turnovers, and recent results from these studies will be presented and discussed.

doi:[10.1016/j.bbabio.2012.06.275](https://doi.org/10.1016/j.bbabio.2012.06.275)

14L2

Functions of the hydrophilic channels of mitochondrial forms of cytochrome oxidase from studies of yeast mutants

P.R. Rich, A. Maréchal, B. Meunier, R. Dodia
 Glynn Laboratory of Bioenergetics, Institute of Structural and Molecular Biology, University College London, Gower Street, London WC1E 6BT, UK
 Centre de Génétique Moléculaire du CNRS, UPR 3404, avenue de la Terrasse, 91198 Gif-sur-Yvette Cedex, France
 E-mail: pr@ucl.ac.uk

Cytochrome *c* oxidase (CcO) catalyses electron transfer from cytochrome *c* to molecular oxygen in a manner that is coupled to

transmembrane proton transfer. Mammalian mitochondrial CcO is a dimer with 13 different polypeptides in each monomer. Subunits I, II and III form its catalytic core and structurally similar homologues of these three subunits are present in the simpler 3–4 subunit bacterial CcOs. The roles of the additional non-core supernumerary subunits of mammalian CcOs are not well understood. Furthermore, despite the structural similarities of the mammalian and bacterial core subunits, difficulties in reconciling existing experimental data preclude a definitive resolution of whether their coupling mechanisms are the same. In particular, a hydrophilic H channel, first identified in the bovine CcO structure [1], has been proposed to be the route for its translocated protons. This contrasts with the view, supported by extensive mutagenesis work, that the D channel fulfils this function in bacterial CcOs. However, an alternative possibility is that the H channel, which is weaker and discontinuous in bacterial CcOs, is functionally required in all forms of CcO but acts instead as a 'dielectric channel' [2] to enable energetically facile transient electron transfer through buried haem *a*. If it does indeed act in this manner, it is further possible that this structure in mitochondrial CcOs may provide, in conjunction with nearby supernumerary subunits, a means of control of core enzymatic function. Evidence for these proposals will be reviewed, together with the new data being obtained with mutant forms of yeast (*Saccharomyces cerevisiae*) CcO [3].

References

- [1] K. Shimokata, Y. Katayama, H. Murayama, M. Suematsu, T. Tsukihara, K. Muramoto, H. Aoyama, S. Yoshikawa, H. Shimada, The proton pumping pathway of bovine heart cytochrome *c* oxidase, *Proc. Natl. Acad. Sci. U. S. A.* 104 (2007) 4200–4205.
- [2] P.R. Rich, Electron Transfer Complexes Coupled to Ion Translocation, In: D.S. Bendall (Ed.), 1996, pp. 217–248, Oxford, UK.
- [3] B. Meunier, A. Maréchal, P.R. Rich, Construction of histidine-tagged yeast mitochondrial cytochrome *c* oxidase for facile purification of mutant forms, *Biochem. J.* 444 (2012) 199–204.

doi:[10.1016/j.bbabio.2012.06.276](https://doi.org/10.1016/j.bbabio.2012.06.276)

14L3

Crystal structure of the alternative oxidases: New insights into the catalytic cycle

T. Shiba¹, Y. Kido¹, K. Sakamoto¹, D.K. Inaoka¹, E. Oluwadare Balogun¹, T. Nara¹, T. Aoki¹, T. Honma¹, A. Tanaka¹, M. Inoue¹, S. Matsuoka¹, H. Saimoto¹, S. Harada², K. Kita¹, A.L. Moore³